

# MANAGING SHEEP IN CONTAINMENT AREAS AFTER FIRE



---

# CONTENTS

<b>INTRODUCTION</b>	<b>3</b>
Welfare requirements	4
Site selection recommendations	4
Construction options	7
Design	9
Farm profile	12
Death rates and culling rates	14
Key recommendations	15
Feeding roughage	19
Animal health	20
How does containment feeding affect wool quality?	22
Release of sheep after containment feeding	23
<b>CASE STUDIES</b>	
Site selection	6
Construction	8
Design	12
Containment area management	16
Mob size	18
Animal health	21

# MANAGING SHEEP IN CONTAINMENT AREAS AFTER FIRE

## A BEST PRACTICE GUIDE

### INTRODUCTION

The information in this publication is adapted from “Managing sheep in droughtlots – a best practice guide”, published by AWI. The practice of containment area feeding is commonly referred to under one of the following terms: sacrifice area, containment area or droughtlot. Containment feeding refers to “maintenance” feeding sheep in contained areas and it can be a valuable tool to managing sheep after a fire. The benefits for stock fed in containment areas after a fire are:

- low stock stress
- stock safe from tree fall
- human safety
- monitoring for maintenance targets
- monitoring for stock health and treating injured stock
- managing feed and water intake

- managing feed and water quality
- assess stock for sale decisions
- manage biosecurity risk from imported weeds in feed
- manage biosecurity risk from wandering stock where boundary fences are breached
- restrict stock access to early colonising plants that affect stock health
- allow pasture recovery
- limit erosion.

Containment feeding can provide a financial saving over paddock feeding, as sheep maintenance requirements are reduced when they are not expending energy walking in search of feed. Table 1 shows over a 9% saving in maintenance energy requirement for medium frame dry ewes of condition score 3 that are containment fed [Source: Lifetime Wool].

**TABLE 1:** Energy required by ewes in condition score 3 (source: adapted from Lifetime Wool).

MAINTENANCE ENERGY (MJ/D) FOR EWES UNDER DROUGHT PADDOCK CONDITIONS							CONTAINMENT FED	
DAY OF PREGNANCY	SMALL FRAME (45KG) MAINTAIN @ CS 3		MEDIUM FRAME (50KG) MAINTAIN @ CS 3		LARGE FRAME (60KG) MAINTAIN @ CS 3		MEDIUM FRAME MAINTAIN @ CS 3	
	SINGLE	TWIN	SINGLE	TWIN	SINGLE	TWIN	SINGLE	TWIN
dry	7.4	7.4	8.0	8.0	9.3	9.3	6.7	6.7
50	7.6	7.8	8.4	8.6	9.7	9.9	7.0	7.2
70	8.0	8.4	8.7	9.1	10.1	10.7	7.4	7.9
100	9.0	10.2	9.9	11.1	11.5	12.9	8.6	9.8
130	11.3	14.1	12.3	15.4	14.4	17.7	10.9	14.1
DAYS LACTATING	MAINTAIN @ CS 3		MAINTAIN @ CS 3		MAINTAIN @ CS 3		EWES AND LAMBS	
	SINGLE	TWIN	SINGLE	TWIN	SINGLE	TWIN	SINGLE	TWIN
10	17.3	21.7	18.7	23.4	21.5	26.9	ask for advice on confinement feeding ewes and lambs	
30	18.7	23.9	20.2	25.8	23.2	29.6		
50	15.5	19.1	16.7	20.6	19.2	23.7		



Containment area refers to the maintenance feeding of sheep in confined areas.

Factors to consider are:

- Drainage on and off.
- Shelter and safety of stock.
- Convenience to facilities.
- Reliable access to adequate quality water supply.
- Minimum distance from water storages and water courses.

The alternative to establishing a specific facility is to use a 'sacrifice' paddock. This can be a degraded pasture paddock that is scheduled to be grazed or renovated. Alternatively, it could be a stubble paddock which has the advantage of a yearly supply of roughage and straw which is useful in preventing soil loss from either wind or water.

The disadvantage of using existing paddocks is that the whole paddock can become bare and subject to damage versus a small area affected if sheep are more confined. Also if mob sizes are not to get too large, for example if weaners and ewes need to be kept separate, a number of paddocks may need to be sacrificed which again increases the area of potential damage.


## WELFARE REQUIREMENTS

The recommendations for sheep welfare are found in the Australian Animal Welfare for Sheep Standards and Guidelines 2016. This document should be read as part of planning noting Chapter 3 Risk management of extreme weather, natural disasters, disease, injury and predation and Chapter 4 Facilities and equipment.

## SITE SELECTION RECOMMENDATIONS

Establishment of a containment area need not be an expensive exercise. However, if possible aim to use existing facilities or if additional facilities need to be constructed try to ensure that they can be useful in the future. On most farms the containment area should be able to be incorporated into existing infrastructure to reduce the cost.

**FIGURE 1:** Possible pen layout option with slope to allow for run off



---

## **DRAINAGE**

Sloping stable soil is preferred to level ground to allow run off, ensure vehicle access to the containment area after periods of rain and to avoid bogging (Fig 1). Site selection after a fire must consider the higher risk of contaminated runoff through the containment area and the risk of tree fall.

## **SHELTER**

Shelter should be considered and if available it should be incorporated into containment areas. Any trees that sheep may have access to will need to be protected to avoid ringbarking. The risk of tree fall after fire and the risk of a second fire event also need to be assessed.

## **CONVENIENCE TO FACILITIES**

Considering the amount of time required to feed, clean and monitor, it is important to make the containment area as close as possible to essential facilities, including fodder storage and sheep handling facilities.



Trees that sheep have access to will need to be protected to avoid ringbarking

## **ACCESS TO A RELIABLE SUPPLY OF ADEQUATE WATER**

Watering from dams is largely discouraged due to the risk of the water supply drying up or becoming contaminated (with soil and dung) following heavy rain. Troughs are generally the preferred option. A large amount of trough space to provide simultaneous access for a large number of sheep is not necessary. Sheep will take turns drinking; high flow rates are more important to ensure rapid replacement of water levels.

## **MINIMAL DISTANCE FROM WATER STORAGE AND WATER COURSES**

A minimum distance of 500m from water storages and water courses is recommended to avoid contamination. Alternatively, a nutrient filter can be located on the down slope side of the area. Contour banks can assist above and below the containment area to minimise water running on and off the site.

## **PRIVACY**

Locate containment area away from houses and public roads.

## **SITE SELECTION CASE STUDY – SEE PAGE 6**

An overview of considerations for site selection as employed by one farm operation in the Southern Tablelands region of New South Wales is provided in Case study 1. The sheep producers in this instance had experience with containment area over two consecutive drought periods. The site selected was intended to support approximately 3,500 mixed age ewes and 1,000 wether weaners.

# SITE SELECTION

## CASE STUDY 1

Case study 1 provides an overview of considerations for site selection as employed by one farm operation in the Southern Tablelands region of New South Wales. The sheep producers in this instance had experience with containment area over two consecutive drought periods. The site selected was intended to support approximately 3,500 mixed age ewes and 1,000 wether weaners.

### FARM PROFILE

<b>Location:</b>	Southern Tablelands NSW
<b>Area:</b>	1,200 hectares
<b>Long term rainfall:</b>	625mm (225mm in 2002)
<b>Pastures:</b>	Mixture of improved perennials, annuals and some native pastures
<b>Enterprises:</b>	Self replacing Merino flock, selling surplus sheep and Dorset x lambs
<b>Management:</b>	August lambing November shearing


The containment area was based on yards built around existing holding yards, a set of sheep yards and a shearing shed. The associated cost of the facilities was low because they were incorporated into existing infrastructure.

### FEATURES OF SITE

- Existing fence lines can be used for one or more sides to minimise costs.
- Ring lock or hinge joint held up by steel posts is commonly recommended.
- Whether the facility is a temporary or permanent fixture can influence the quality of the construction.

### FACILITY

CASE STUDY 1 - Pen layout based around existing sheep yards



---

## CONSTRUCTION OPTIONS

Recommendations for construction:

- Existing fence lines can be used for one or more sides to minimise costs.
- Ring lock or hinge joint held up by steel posts is commonly recommended.
- Whether the facility is a temporary or permanent fixture can influence the quality of the construction.
- Merinos do not require fencing of the standard used for normal farm fences, which helps to minimise cost. More robust facilities are required for British breeds and their crosses.

## CONSTRUCTION CASE STUDY – SEE PAGE 8

An example of construction methods undertaken by one farm in the Southern Tablelands region of New South Wales is provided in Case study 2. The sheep producers in this example had no previous experience in establishing a containment area facility. The aims of this particular containment area were to:

- Keep options open.
- Protect the land.
- Maintain stock numbers after the drought.



Existing fence lines can be used for one or more sides to minimise costs.

---

# CONSTRUCTION

## CASE STUDY 2

Case study 2 provides an example of construction methods undertaken by one farm in the Southern Tablelands region of New South Wales. The sheep producers in this example had no previous experience in establishing a containment area facility. The aims of this particular containment area were to keep options open, to protect the land and to maintain stock numbers after the drought.

### FARM PROFILE

<b>Rainfall:</b>	Average 500mm
<b>Enterprises:</b>	Merino flock and cattle
<b>Management:</b>	October shearing Mid June lambing

### CONSTRUCTION

The containment area consisted of four holding pens 75m wide by 100m deep. In front of these were two feeding pens each 30m x 230m. It took approximately one month to construct the containment area to hold 6,000 sheep.



Existing fence lines can be used for one or more sides to minimise costs.



Recycling or using existing infrastructure is recommended.

Feeding pens were constructed separate to holding pens.

Fences were constructed of creosote posts at 10m spacings and a dropper in between. Sheep and lamb cyclone was used on all except the high pressure areas. Around the outside of the containment area a plain wire was added on top. High pressure areas had 1m high pig cyclone (approx 15cm mesh). Gates were 2 x 10m for each pen.

A handling facility was built in the corner of one of the feeding pens.

### CONSTRUCTION

- Cost of materials was \$10,000.
- Materials included one water tank and four troughs, posts and cyclone.
- Labour to install was around \$5,000 which equates to \$4.50 per sheep.
- The facility will be used again.

## DESIGN

There is no standard for design of containment areas, however the three most commonly used designs feature on pages 9-11. The advantages and pitfalls of each design are specified.

### OPTION 1: SEPARATE FEED YARD

Hay is fed in the holding yard while grain is fed in the feeding yard. Sheep are kept in the holding yard except on feeding days (Figure 2).


## PROS

- Avoids the need for troughing for each mob.
- Grain can be fed out in a pen without sheep being present, which makes it easier than driving through the mob of sheep.

## CONS

- Requires feeding of hay and grain to at least one mob every day, even if each mob is fed 1-2 times per week.
- Additional fencing and gates compared to Option 3.
- Increased risk of diseases associated with contamination of site, e.g., salmonella, coccidiosis.

FIGURE 2: Separate feed yards



Source: Feeding Sheep, PIRSA (1997)

---

## OPTION 2: EXTRA PEN ROTATION

The design is based on having one more pen than mobs of sheep (Figure 3). This enables sheep to be fed in empty pens and then stock let in. Grain is fed first in the empty pen then stock moved in. Feed is then put out in the next pen and stock moved in and so on. At the next feed the process is reversed with feed put out without stock present in the pen.


### PROS

- Feed can be put out in empty pen.
- All yards can be fed on the same day.

### CONS

- Feed and water troughs required in each pen increases cost.
- One more pen required than number of mobs being fed.
- If pens are small it does not enable one length of troughing when following standard recommendations of 15-20cm of double sided trough per 100 head.
- Sheep rushing through gates to feed can be a problem. They appear to improve with time.

FIGURE 3: Extra pen rotation



---

### OPTION 3: ONE YARD PER MOB

This design provides one yard per mob to be contained (Figure 4). The capital cost is minimised but feeding can be more difficult unless fed from the outside of the yard.


#### PROS

- Minimise fence length and therefore cost.
- All yards can be fed on the same day.
- Troughing outside the yard makes feeding easy and a one person job.
- Troughs outside the yard minimise the risk of disease due to site contamination, e.g., salmonella, coccidiosis.

#### CONS

- Feed is put out while sheep are in pens, increasing the chance of sheep escaping or moving between pens.
- May require two people to feed out.
- Troughs outside the yard doubles the length of trough required because sheep can only get access to one side. Also requires a system that delivers feed from the side of vehicles.
- Small pen sizes do not enable one length of troughing in the pen when following standard recommendations of 15-20cm of double side trough per 100 head.

FIGURE 4: One yard per mob



# CONSTRUCTION

## CASE STUDY 3

An overview of the design of a containment area in southern NSW is provided in Case study 3. The sheep producers in this instance were experienced in containment area management having used them previously, in the droughts of 1994 and 1997. Important lessons in design from the earlier years were transferred and expanded upon to result in a highly functional and practical lot system.


### FARM PROFILE

<b>Area:</b>	1,880 hectares in four different blocks
<b>Rainfall:</b>	Long term average 800mm (587mm in 2002)
<b>Pastures:</b>	80 per cent of country sown down to perennials
<b>Enterprise:</b>	Self replacing fine wool (18.5 micron) Merino flock of 12,000 ewes, 8,000 wethers and 4,500 weaners

### CONTAINMENT AREA DESIGN

- Aim to stock at 5,000 – 6,000 per hectare.
- To avoid opening gates and minimise feeding time, always feed from the outside of the containment area (using a side delivery system).
- Gates should be on the opposite side of the pen to the feed troughs to make removal of shy feeders easier.
- Construct pens as part of existing facilities wherever possible to ensure they can be used at other times.

### CASE STUDY 3: Fence and feed trough system



---

# CASE STUDY 3 (CONTINUED)

## FEED TROUGHS

Provide 5cm of trough space per head.

Trough capacity must be at least 1.5kg per head. Troughs were rectangular running down the long side of the pen.

Locate feed troughs on the top side of the containment area so run off moves away from the trough. A number of different types of feed trough were used, including:

- Conveyer belt with fence in the middle. The main problem was sheep could stand in them so they needed regular cleaning but they were also slow and therefore expensive to construct. Belting cost increased to \$12 per metre during the drought.
- To overcome the problem of contaminating the troughs, belting was tied on the outside of the fence with the fence high enough off the ground to allow sheep to feed underneath. This was not satisfactory because the belting was too wide and sheep could not reach all the grain. Also there was the occasional tyre spiked on the cut off steel posts used to support the belting.
- The third and successful option was to use metal feed troughs on the outside of the fence with a 175mm gap between the top of the trough and bottom of the fence. Troughs cost approximately \$11 per linear metre and will need to be stored inside when not used. They are much easier to handle than belting.



Feeding from outside pens requires a side delivery system.  
©CSIRO Plant Industry



Provide 5cm of trough space per head.  
©CSIRO Plant Industry

- Provide 5cm of trough space per head. Provision of at least 5cm of trough space per head was obviously insufficient for all sheep to feed at once but all seemed to have sufficient access. Those mobs with more trough space had no more tail than those with the minimum 5cm.

## LESSONS LEARNED AND THINGS TO CHANGE

- Incorporate the facilities into the design of your existing yards wherever possible.
- Ensure the water system is right from the start.
- Water supply must be good – it is easier to move the sheep and the grain to water than vice versa.
- Feed lime from small troughs in the pens, not through the feed-out cart because it affects the flow rate of the grain.
- Locate feed troughs on the top side of the containment area so run off moves away from the trough.

## DEATH RATES AND CULLING RATES

There is a wide variation in death rates in containment area. Surveys in South Australia showed an average of 1.8 per cent and 1.4 per cent in 1982 and 1988 respectively. An AWI survey found an average of 2.8 per cent and a median of 1.4 per cent. The distribution of deaths is shown in Table 2.

The main point to note from these results is that the majority of sheep fed in containment had low levels of mortality and much lower than the potential death rates if sheep were not supplementary fed during drought. However, there were a few which experienced high death rates for a variety of reasons.

Table 3 outlines causes of mortality as recorded in the AWI-commissioned survey. The data reveals acidosis and tail end sheep (responsible for 85 per cent of total deaths in containment area during the 2002/03 drought period) as the more important causes of death.

**TABLE 2:** Percentage of farms by death rate

DEATHS	1982	1988	2002/03
0-1%	60%	65%	70%
1-2%	18%	10%	16%
2-3%	8%	10%	4%
3-4%	5%	10%	2%
>4%	11%	5%	8%
	(Ashton & Hannay 1984)	(Morbey & Ashton 1990)	(AWI/HSA 2003)

**TABLE 3:** Causes of mortality

CAUSE	%
ACIDOSIS	48
TAIL END*	37
ACCIDENTAL	3
UNKNOWN	16

\* Poor condition

Source: Holmes Sackett & Associates

Implement a number of management practices to minimise the risk of deaths, including:

- Introducing grain according to recommended schedules prior to introduction to the containment area if possible.
- Removing shy feeders prior to entry and whilst in the containment area. If there are high stocking densities or if there is a tail appearing in the mob, this should be done at least every two months and more frequently in larger mobs.
- Feeding roughage.
- Ensuring sheep are properly vaccinated against clostridial diseases. Seek veterinary advice to assess if your annual vaccination regime for clostridial disease is frequent enough to protect against the higher risk of pulpy kidney when feeding in containment areas.
- If possible avoid running young sheep in containment area.
- Sheep rushing through gates to feed can be a problem. They appear to improve with time.

## KEY RECOMMENDATIONS

Following the 2002/03 drought, AWI commissioned a survey of sheep producers who managed sheep in containment areas. The survey was undertaken to collect some basic information on the performance of sheep in containment areas and to identify any factors that were an important influence on the survival of sheep.

Participants were predominantly from New South Wales and Victoria with a small number from South Australia. A total of fifty sheep producers were involved in the survey and, allowing for the fact that many had more than one management group, a total of 125 different containment areas were surveyed.

The average and range of results is set out in Table 4.

Based on the results of the analysis there were a number of factors that were shown to influence the performance of sheep in containment areas. These factors provide a basis for setting up and managing sheep in confined areas but they are a guide only. In a number of instances the management program can be varied considerably. This applies particularly to mob sizes.

## STOCKING DENSITY

Because pasture availability on a partially burnt property may be low until significant rainfall, containment areas can be needed for an extended period of time and careful selection of sheep to be fed in these areas is important. The recommended



Higher stocking densities than 2,000 sheep per hectare tended to be associated with higher mortality rates and poor doers.

**TABLE 4:** Average and range of results from AWI containment area survey 2002/03

	AVERAGE	RANGE
TIME ON FEED (DAYS)	136	41-407
STOCKING DENSITY (SHEEP/HECTARE)	1,210	2-9,862
MOB SIZE (NUMBER OF SHEEP)	1,214	22-12,222
SUPPLEMENTARY FEEDING PRIOR TO ENTRY (WEEKS)	6.8	0-40

## CONTAINMENT AREA MANAGEMENT CASE STUDY – SEE PAGE 16

An example of containment area management methods adopted on one farm located in central New South Wales is provided in Case study 4. This example highlights the importance of creating a plan of action and schedule based on key strategies for selling and feeding.

stocking density is approximately 2,000 sheep per hectare. In the survey, higher stocking densities tended to be associated with higher mortality rates and poor doers. This is particularly important if roughage is not going to be fed, because high stocking densities without roughage are associated with a substantial increase in poor performance. Very low stocking densities also tended to be associated with poor sheep performance, possibly because of inadequate access to feed. Low stocking densities also resulted in more country being damaged, which to some extent defeats the purpose of one key benefit of a containment area.

Many mobs have been run at stocking densities well over 2,000 sheep per hectare with one of the objectives being to minimise dust. At high stocking densities of around 5,000 sheep per hectare, the urine and dung may be more likely to form a hard pad, which minimises the problem with dust and even mud. Whether or not a hard pad forms seems to be more dependent on soil type than stocking density with some soils not packing hard, regardless of stocking density.

One advantage of higher stock densities is that less infrastructure is required for a given number of sheep.

# CONTAINMENT AREA MANAGEMENT CASE STUDY 4

Case study 4 provides an example of containment area management methods adopted on one farm located in central New South Wales. This example highlights the importance of creating a plan of action and schedule based on key strategies for selling and feeding.

## FARM PROFILE

<b>Location:</b>	Central NSW
<b>Area:</b>	1,400 hectares
<b>Rainfall:</b>	Long term average 800mm
<b>Pastures:</b>	1,000 hectares sown to improved pastures
<b>Enterprise:</b>	7,000 Merino sheep and a 500 head self replacing beef herd

## STRATEGY

The most important factor in the whole feeding program was writing down the sell or feed strategy. The sheep producers wrote down dates at which time key actions were to be undertaken if rain had failed to arrive in spring 2002. These dates were cast in stone.

DATE	ACTION
1 August	Ring the stock agent and sell 1,350 older wethers.
1 September	Draft off any cull ewes and sell. Accept the market price. Plan feed requirements for the next 8 months and consider locking in a feed grain contract. Assess cattle feed available and sell any trade animals that are ready. Do not be greedy.
1 October	Start introducing wheat rations. Start building feed pens. Check on feeding equipment (augers, silage carts etc). Have all feed on hand tested for quality.
1 November	Start moving sheep into pens.

By acting early, a good price was obtained for sale wethers and cull ewes. While target dates slipped a bit, sheep were taken off the paddocks before any serious damage was done to pastures. The program was not deferred just because rain looked likely at the time.

Lambs were weaned earlier than usual. They were all trained to feed on grain while on their mothers, and were fed a lupins/wheat mix.

The tail of each mob was drafted off before containment area entry and then regularly during lot feeding.

The advantages achieved from lot feeding that are considered important were:

The preservation of pastures. Having undertaken extensive pasture improvement over the past eight years the sheep producer wished to protect the capital investment.

No plan was ever made to containment feed lambs because they had many more paddocks available on which to be grazed.

Ease of feeding stock. It is quicker and more efficient to feed stock in pens.

Shy feeders were able to be identified, easily separated and fed preferentially.

## MOB SIZE

The optimum mob size is less than 1,000 head. There is no benefit in having smaller mobs, so mob sizes up to 1,000 should be determined more by logistics (e.g., number in age groups or classes). Mob size can be increased if necessary but it will be likely that additional management will be required to identify and remove tail end sheep. Mob sizes should not exceed 2,000 head because it presents an increased risk of progressively higher mortality and culling rates particularly when sheep are fed on the ground. If large mob sizes are needed, it is preferable to feed sheep in troughs (Graph 1).


### MOB SIZE CASE STUDY - SEE PAGE 18

An example of feeding practices, costs and lot designs for a large number of smaller mobs located on a farm in central Victoria is provided in Case study 5. The practice of feeding the mobs on the ground rather than in troughs worked well in this instance.



The tail end of the mob should be removed at least every two months ©CSIRO Plant Industry.

GRAPH 1: Interaction between mob size and mortality



## TAIL MANAGEMENT

The tail end of the mob should be removed at least every two months, particularly if mob sizes are in the range of 1,000 to 2,000 head. Failing to do this will increase the risk of deaths. Poor doers may need removing more frequently in the early stages, though these may not be very obvious if sheep are in good condition on entry. An area designed like that in Figure 1 allows easy identification and removal of shy feeders because they do not move across to the feed.

Factors which contribute to a large tail in the mob include:

- Feeding grain only without roughage (see separate section on roughage).
- High stocking densities (greater than 2,000 sheep per hectare).
- Very low stocking densities.
- Mob sizes which exceed 1,000 head.

If you have any of these factors in the containment area be prepared to remove the tail of the mob more frequently to minimise the number of tail end sheep that die.



---

Removing the tail end can be easily done by eye through the draft process. There is no need to weigh or condition score sheep to do this well because the poor doers are obvious, particularly if in short wool.

Weighing and/or condition scoring a sample of the ewes is useful for monitoring mob performance and provides a means of adjusting rations when required.


### CLASS OF SHEEP

It is usually easier to manage adult sheep rather than weaners in containment areas, so if possible leave the weaners out on pasture. If it is necessary to confine weaners it can be done, but they are likely to need additional management. Weaners are also more susceptible to problems associated with disease.

### TROUGHS

Troughs are not essential for successful management of sheep in containment areas. Feeding directly onto the ground actually reduces the risk of deaths associated with smothering and poor doers in the mob (Graph 2).

**GRAPH 2:** The effect of troughs on mortality rate



One of the main advantages provided by troughs is to minimise feed wastage in wet weather so if not using troughs ensure you have a strategy to manage wet conditions. The most obvious way is to let sheep out into a paddock for several days and feed them there until the containment area dries out.


### FEEDING ROUGHAGE

Many departmental publications provide comprehensive information on the feed requirements of sheep and these should be referred to when calculating quantities to feed.

Sheep producer experience, trial work and analysis of survey results show that there is a benefit in including some roughage in the ration. The AWI survey of the 2002/03 drought showed that feeding roughage (hay, straw or silage) reduced death rates from an average of 1.34 per cent to 0.76 per cent (Graph 3).

During previous droughts in South Australia, research showed that mobs fed grain only had no higher mortality rate but did have a greater proportion of sheep removed as poor doers.

**GRAPH 3:** Effect of roughage in diet on mortality



In addition the work showed that those mobs fed straw rather than hay had fewer poor doers (Table 6). Roughage should be fed at a minimum of 1kg/head/week.

Roughage does not need to be high quality and if only small quantities of roughage are to be fed, it is best to feed low quality roughage such as straw. This helps to ensure that all sheep in the mob have access to some roughage whereas high quality roughage is consumed rapidly by a small number of aggressive feeders.

### ANIMAL HEALTH

In addition the work showed that those mobs fed straw rather than hay had fewer poor doers (Table 5). Roughage should be fed at a minimum of 1kg/head/week.

- Acidosis associated with high grain intakes of animals that are not accustomed to such rations. To minimise the risk, ensure animals are introduced to grain over the time recommended in state department publications. The schedule usually requires 3-4 weeks to build the ration up to that required for maintenance feeding. Some sheep producers experienced problems with acidosis towards the end of the scheduled introduction period when daily rations were increased from 430g/head/day up to 860g every second day. Producer experience has shown that there is less risk of acidosis if this increase is done with an extra step when sheep are fed 650g per head every second day.



Animal health problems in containment areas tend to be concentrated on a few factors which can cause substantial health problems if not managed well. ©CSIRO Plant Industry

**TABLE 5:** Inclusion of roughage

	MINIMAL ROUGHAGE		MODERATE ROUGHAGE	
	HAY	STRAW	HAY	STRAW
QUANTITY (KG)	0.5	0.7	1.0	1.4
POOR DOERS (%)	25	8	18	3

Source: Brian Ashton and Alex King, PIRSA

- Access to roughage is also important if sheep are being introduced to grain in the containment area rather than at pasture. If possible, sheep should always be introduced to full grain rations prior to introduction to the containment area to minimise the risk of acidosis and to identify as many shy feeders as possible.
- High levels of grain feeding predispose sheep to enterotoxaemia (pulpy kidney). To minimise this risk, sheep should have at least two vaccinations at least one month apart with the second at least two weeks prior to commencement of grain feeding. Young sheep are the most likely not to have had a complete vaccination history. Older sheep are likely to have had a number of clostridial vaccinations over their life so are more likely to have a higher level of immunity. A booster should still be given to overcome the higher risk that long periods of grain feeding present.
- Internal parasites can be a problem in containment areas simply due to the high concentration of sheep. A drench prior to entry is worthwhile and worm burdens should be monitored by use of faecal egg counts, particularly in young sheep. Young sheep tend to present a higher risk.
- Some diseases are more likely in concentrated mobs. The two main risks are coccidiosis and salmonella infection in stressed or young sheep. Contaminated feed and water can be predisposing factors. If you have concerns at any stage, investigate problems as soon as they start to appear because high concentrations of sheep will encourage a rapid spread of any problems. Consult a veterinarian.

---

# ANIMAL HEALTH

## CASE STUDY 6

Case study 6 focuses largely on nutritional concerns. It outlines how one sheep producer managed problems with infection which arose in a mob of weaners during containment feeding.

### FARM PROFILE

<b>Area:</b>	1,417 hectares
<b>Pastures:</b>	90 per cent pasture improved (rye, sub-clover, phalaris, cocksfoot, fescues)  Maintenance level super applied to 100 per cent pasture each year for last 20 years
<b>Enterprise:</b>	Self replacing Merino flock

### MANAGEMENT

Once the weaners had been introduced to the grain ration, they were moved into the first intensive containment area yard. Maiden ewes were still in the paddock and maintaining weight.

- The weaners were maintaining condition and consuming 350g per day of the ration.
- The maiden ewes had a condition score of 2.5-3.0 across the flock.



The recommended three drenches for coccidiosis were administered on three successive days.

Lambs were looking weak and began dying overnight. Given the rapid onset of the problem a solution had to be found quickly. A local vet confirmed coccidiosis and recommended that three drenches be administered on three successive days.

The infection was halted after intensive drenching.

There was a second outbreak 3-4 weeks later. To combat the second outbreak sheep were treated using a drench and move system at 3-5 day intervals. This program proved as successful as the one recommended by the vet.

The coccidiosis was attributed to stress at weaning, feeding off the ground in the introduction paddock and the intensity in the containment area situation. Having the ability to rotate through several pens was important to improve hygiene when sheep, particularly young sheep, are confined for long periods.

Worms were monitored closely for both mobs through the whole exercise. They indicated that no drench was necessary for the weaners or the maidens through the entire confinement.

The computer program GrazFeed, a decision support tool that helps calculate sheep and cattle feed requirements, was used to formulate the rations. It recommended a mix of 70% oats, 20% lupins, 10% hay and 1.0%-1.5% lime for the lambs. This proved to be accurate as the weaner condition score did not vary much over the five months but they did grow steadily. The maidens were fed the same mix as the weaners but adjusted up for their liveweight. This saved time on preparing rations.

Average feed cost per week for the weaners was \$0.87 and \$0.92 per week for the ewes.

Feeds were sourced on price and quality. Wheat and lucerne hay were available and used depending on value. Grains were kept more constant due to difficulty and inefficiency in random changes. Over time, various grains were blended into the ration including triticale, barley, and sheep nuts.

---

# CASE STUDY 6 (CONTINUED)

## DEATHS

Some lambs were lost due to smothering whilst trying to access hay as it was deposited in the pen. Total lamb deaths were estimated at 7 per cent. Lambs were also lost in the pen with the dam in it.

Ewe mortality was low (<1 per cent) as they did not experience any significant disease problems. Few experienced any grain poisoning or feed-related complications.

## LESSONS LEARNT AND THINGS TO DO DIFFERENTLY NEXT TIME

Using a specialist nutritionist would be something considered next time. Greater use of specialist knowledge in sheep requirements and finetuning the ration would have saved money.

## HOW DOES CONTAINMENT FEEDING AFFECT WOOL QUALITY?

Containing sheep at high stocking rates in small areas can have a number of effects on wool quality. The experience of sheep producers has shown that there is considerable variation in the effect of containment areas on wool quality. The main issues have been:

- Management of staple strength, particularly as sheep are introduced and released from the containment area. The key is to manage the change in nutrition as smoothly as possible. Moving sheep from the paddock to the containment area should be done as smoothly as possible with a gradual introduction of grain. Conversely, when releasing sheep, going from a maintenance ration in the containment area to high quality pastures, with for example 1,500kg dry matter/hectare, will result in rapid weight gain and a rapid change in the fibre diameter profile. Rapid changes in fibre profiles will result in reduced staple strength. The aim should be to ensure sheep go from a maintenance containment area ration to a maintenance pasture ration. Any increase above maintenance

should be as gradual as possible.

- Low staple strength is not an inherent problem associated with containment areas, rather it is about how sheep are managed into and out of the containment area that is important.
- High dust levels in some mobs. This was a particular problem with sheep going into containment areas with long wool. Yields of 50 per cent or less were common in these situations. Extremely low yields will affect the clean price of wool, particularly when there is an increase in supply. Wherever possible, it is preferable to have sheep in short wool while they are confined. This is not always going to be possible because a premature shearing may be required which in turn may result in larger discounts than the dust in full length wool. Consider your options and do your sums carefully.



Containing sheep at high stocking rates in small areas can have a number of effects on wool quality.

---

## RELEASE OF SHEEP AFTER CONTAINMENT FEEDING

Weaning sheep out of the containment area and back into the paddock is considered important to avoid any marked break in the wool. Some sheep producers have found it useful in the past to let sheep out onto feed for a few hours each day after feeding over a set period of weeks.

To reduce the likelihood and risk of digestive disturbances, it is recommended that sheep are released from containment area facilities after feeding. It is also necessary to continue with drought rations until pasture dry matter has reached sufficient quantities to meet the requirements of the sheep.

Late pregnant or lactating ewes should continue to be fed limestone and salt mix.

Delaying release will increase dry matter (DM) availability and provide optimum growth rates if pastures reach approximately 1,500kg of DM/hectare. In many situations, this is unlikely to be economic because at a growth rate of 50kg/



Weaning sheep out of the containment area and back into the paddock is considered important to avoid any marked break in the wool.

hectare/day after the drought breaks, it will take 30 days to reach 1,500kg assuming zero decay.

In some instances, the cost of maintaining sheep in the containment area will mean sheep are released well before pastures reach 1,500kg. Dry sheep being fed a maintenance ration in a containment area will continue to maintain liveweight on pasture with 400-500kg of DM/hectare.

AWI's publication 'Releasing sheep from containment feeding' provides a lot of detail around important considerations for releasing sheep from containment feeding, including animal health and minimum pasture requirements.

## SHEEP PRODUCER COMMENTS

Of those surveyed, 98 per cent said that they would use a containment area again if the situation demanded it.


Other general sheep producer comments included:

- Weaners are more difficult to manage.
- Roughage appeared to be important.
- Dust is a problem.
- Do not need purpose built facility.
- Hard on stock and not a good experience but prepared to do it again because the alternative is worse.

This publication has been printed on ecoStar paper



AWI has a range of planning, management and recovery resources available for sheep producers going into, enduring and recovering from drought which are also applicable to recovery from fire. For your free copy, visit [www.wool.com/drought](http://www.wool.com/drought) and [www.wool.com/land/bushfires](http://www.wool.com/land/bushfires) or call the AWI Helpline on 1800 070 099.



### **Managing fodder prices for drought:**

A guide which focuses on strategies to help sheep producers manage fodder prices and supply risks during droughts.



### **Which sheep do I keep?**

A guide to help sheep producers decide which sheep to keep during drought. The guide helps managers, when confronted with a pasture shortage, determine whether to sell or supplementary feed all, some or none of their flock.



### **Stock water - a limited resource after fire:**

Calculate stock water budgets - critical information for any sheep producers considering containment feeding their sheep.



### **Releasing sheep from containment feeding**

It's rained and you're preparing to release your sheep from containment. Before you do, take a look at this guide to understand the minimum pasture levels and key animal health considerations for a successful release.



**WOOL.COM**