



IT'S FLY TIME! RECOGNISING AND MONITORING FLYSTRIKE

KEY POINTS

- Australian sheep blowflies (*Lucilia cuprina*) initiate most flystrike, but other species can cause additional and sometimes more severe damage.
- Aim to detect flystrike as early as possible to prevent severe infections.
- Monitor all mobs of sheep for signs of flystrike during high-risk periods, but especially weaners.
- Be aware of and monitor for weather conditions which may result in increased flystrike risk.
- Monitoring for flystrike in sheep should be included in a broader plan which also considers prevention and treatment activities.

WHAT DO AUSTRALIAN SHEEP BLOWFLIES LOOK LIKE AND WHAT IS THEIR LIFECYCLE?

The Australian sheep blowfly (*Lucilia cuprina*) is responsible for initiating 90% of flystrike in Australia. The adult Australian sheep blowfly is about 9mm long (body length) and metallic green/bronze in colour with reddish eyes.

After the Australian sheep blowfly has initiated a strike, other species of fly may attack the animal and can cause additional damage which is sometimes more severe.



An adult female (left) and male (right) Australian sheep blowfly

Source: J Larsen, L Tyrell and N Andersen, The Mackinnon Project, University of Melbourne, from *Early Season Treatment and the Control of Breech Strike in Unmulesed Sheep – a National R&D Technical Update* (2010), AWI



Various stages of Australian sheep blowfly larvae development

Source: S De Cat and J Larsen, The Mackinnon Project, University of Melbourne, from *Managing Breech Flystrike* (2019), AWI

Warm, moist, humid and calm conditions can lead to damp, smelly wool which is sought after by egg-bearing female flies as an ideal environment for the larvae which will hatch from their small white eggs.

After hatching, larvae need a feed rich in protein to commence their lifecycle. This can be sourced from wounds, existing strikes, weeping skin, carcasses and protein-rich manure.

Larvae (creamy coloured maggots) hatch within 12-24 hours of the eggs being laid, grow by feeding on the sheep through wounds, existing strike and wet (e.g. wet due to urine and dags) or weeping skin.

The maggots drop off the sheep after about three days and burrow into the ground to pupate. The pupa become barrel-shaped as the outside shell hardens and darkens from yellow to red-brown. When conditions are favourable, the young flies emerge about one week later.

HOW DO I KNOW IF AUSTRALIAN SHEEP BLOWFLIES ARE AROUND?

The most obvious sign will be fly struck sheep but there are other indicators to consider.

Consecutive warm, wet days provide ideal conditions for female flies to lay eggs on sheep and are a good indicator that strike may occur. Adult flies do not normally travel further than 3 km and sheep camps and watering points can act as epicentres for infection. Check these areas regularly for blowflies and struck sheep.

Fly traps may be useful tool to monitor fly activity in northern areas of New South Wales and Queensland. If any Australian sheep blowflies are caught in traps, this should be taken as a warning that flies are about and, given favourable conditions, flystrike will occur. In southern regions, they may be useful for determining when flies are emerging from overwintering and the fly season is about to commence.

WHAT ARE THE SIGNS OF FLYSTRIKE?

The aim of all monitoring activities should be to detect and address flystrike as soon as possible.

COVERT FLYSTRIKE

Small areas of flystrike that cannot be detected easily are quite common and are known as covert flystrikes. They are difficult to detect unless you are looking closely at sheep, such as when handling them for crutching, shearing or marking.

Covert flystrike can last for some weeks before advancing into more obvious flystrike during warm, moist conditions or they can resolve without the need for treatment.



Covert flystrike can be difficult to detect unless handling sheep

Source: FlyBoss

EARLY DETECTABLE FLYSTRIKE

Early signs of flystrike which should be noted in a monitoring program include patches of wool that appear discoloured from chewing or rubbing. Early detectable flystrike does not usually have a large strike wound but the discolouration becomes progressively darker with more exudate (weeping).

Sheep also usually behave differently as they will be uncomfortable and irritated. They may twitch their tail, bite or scratch at the affected area if they can reach it, especially with breech strike. They may also stamp their feet, duck their heads and arch their backs but will generally stay with the mob.

As soon as an early detectable flystrike is confirmed, implement treatment activities and take steps to prevent further flystrike from occurring. Further details on the prevention and treatment of flystrike can be found in AWI's It's Fly Time! Preventing and treating flystrike factsheet available from www.wool.com/itsflytime.



Example of early detectable strike

Source: AWI

Identifying why the strike occurred will help you work out whether it is an anomaly (only one sheep is affected), in which case dramatic intervention may not be warranted, or if it is an indicator that an outbreak is imminent. Consider if the strike occurred due to a management error (e.g. poor or delayed crutching, chemical application error, failure to implement preventative action soon enough in a normal season) or due to the susceptibility of your sheep.

If you think an outbreak might happen, increase monitoring and take action such as crutching or chemical preventions as soon as possible. If shearers cannot be arranged, consider alternative options, such as additional jetting which may provide interim protection until shearing or crutching can be arranged.

ADVANCED FLYSTRIKE

The signs of advanced strike are more obvious because the wounds are larger.

Wounds from advanced strike are large, wet and dark, and maggots will be moving outwards to consume healthy tissue.

Sheep will be suffering significant pain as well as fever and other whole of body effects.

Sheep won't be able to keep up with the mob when grazing and are often found on their own. They will become increasingly impacted by the flystrike, stop eating and drinking, will lie down and not want to get up.

Strike from secondary blowfly species can be smaller but more serious as the maggots invade deeper tissue not usually directly impacted by the Australian sheep blowfly, causing extensive pain, tissue damage and illness.

Without treatment, sheep will generally die quickly, anywhere from hours to a few days. If sheep with advanced flystrike are found, treat animals immediately and increase the frequency of monitoring as it is likely that more flystrike will occur in the following days or weeks. Read more about treatment in It's Fly Time! Preventing and treating flystrike.

When monitoring for flystrike, the aim is to detect strike as early as possible to minimise pain and suffering.



Example of advanced strike

Source: AWI

WHAT WEATHER CONDITIONS FAVOUR FLIES?

Blowfly eggs hatch when the temperature is above 15°C with blowfly activity increasing above 17°C and peaking between 26°C to 36°C. The longer the temperature remains above 17°C, the greater the chance of egg laying and dispersal.

Consistent, regular rainfall that keeps the skin of the sheep moist for more than two days increases the risk of flystrike. Such conditions also cause odours from fleece rot, dags and urine stain to be more prevalent and allow skin folds to remain moist, attracting female flies to lay eggs. Frequent, small falls of rain are more conducive to flystrike than occasional heavy falls because there is less opportunity for the skin to dry out.

Fly activity is reduced when wind speeds are greater than 9 km/h and stops when wind speed is greater than 30 km/h.

This combination of weather conditions usually coincides with spring and autumn, but flystrike can occur throughout the year if climatic conditions favour fly reproduction.

Aim to carry out prevention strategies before temperature and rainfall increase and monitor sheep carefully when favourable flystrike conditions occur.

FURTHER INFORMATION



To access the other It's Fly Time!™ tools and information mentioned in this document visit www.wool.com/itsflytime

OTHER INFORMATION



To access a wealth of information on strategic flystrike management visit www.wool.com/simplifly



For more information on the AWI Flystrike Extension Program visit www.wool.com/flystrikeresources



For all the latest flystrike R&D publications in one spot visit www.wool.com/flystrikelatest



To access a wealth of information about breeding and selection for flystrike resistance visit www.wool.com/breedingforflystrikeresistance

FLYBOSS

Lice and flystrike product tool

www.flyboss.com.au/sheep-goats/treatment/products

Flystrike Quick Tools (online)

www.flyboss.com.au/sheep-goats/tools/flystrike-quick-tools-online.php

Flystrike Risk Simulator (downloadable)

www.flyboss.com.au/sheep-goats/tools/flystrike-risk-simulator-downloadable.php

National risk maps for flystrike

www.flyboss.com.au/sheep-goats/management/national-risk-maps

OTHER PUBLICATIONS

Sheep blowflies Primefact 485 (NSW Department of Primary Industries)

www.dpi.nsw.gov.au/_data/assets/pdf_file/0005/289643/Sheep-blowflies.pdf

Managing flystrike in sheep (WA Department of Primary Industries and Regional Development)

www.agric.wa.gov.au/livestock-parasites/managing-flystrike-sheep

FlyBoss Monitoring for flystrike

www.flyboss.com.au/sheep-goats/management/monitoring-for-flystrike.php

The biology and ecology of the Australian sheep blowfly (Wiedemann) – an update (KG Wardaugh, proceedings from the FLICS Conference)

www.licebosstools.org.au/FLICS/PDFs/Chapter_2/Wardaugh_Lucilia_biology_and_ecology.pdf

Factors affecting the incidence of flystrike in sheep – a description and analysis of data from three separate areas in eastern Australia (KG Wardaugh, RJ Mahon and D Bedo, Proceedings from the FLICS Conference)

www.licebosstools.org.au/FLICS/PDFs/Chapter_2/Wardaugh_et al_Factors_affecting_incidence_flystrike.pdf

Published by Australian Wool Innovation Limited, Level 3, 24 York Street, Sydney NSW 2000.

This publication should only be used as a general aid and is not a substitute for specific advice. To the extent permitted by law, we exclude all liability for loss or damage arising from the use of the information in this publication.

© 2022 Australian Wool Innovation Limited. All rights reserved. AWI invests in research, development, innovation and marketing activities along the global supply chain for Australian wool. AWI is grateful for its funding, which is primarily provided by Australian woolgrowers through a wool levy and by the Australian Government which provides a matching contribution for eligible R&D activities. GD4108A_Jul2022